

# Materials Compatibility Reference



## Compatible Materials

**Prepare or administer DOJOLVI using containers, dosing syringes, or measuring cups made of these materials, such as:** High-density polyethylene (HDPE), low-density polyethylene (LDPE), polypropylene, stainless steel, glass, polyurethane, and silicone























## Incompatible Materials

**Do not prepare, administer or store Dojolvi using these materials:** Polystyrene or polyvinyl chloride (PVC)



## Unknown Compatibility

Compatibility for these materials is unknown. **Use is not recommended.**

  <b>PETE</b>	  <b>HDPE</b>	  <b>PVC</b>	  <b>LDPE</b>	  <b>PP</b>	  <b>PS</b>	 <b>Other Compatible Materials</b>
<p><b>polyethylene terephthalate</b></p>	<p><b>high-density polyethylene</b></p>	<p><b>polyvinyl chloride</b></p>	<p><b>low-density polyethylene</b></p>	<p><b>polypropylene</b></p>	<p><b>polystyrene (also known as styrofoam)</b></p>	<p><b>stainless steel, glass, polyurethane, silicone</b></p>
<p>examples include: soft drink bottles, water bottles, and fruit juice containers<sup>1</sup></p> 	<p>examples include: milk, juice or water bottles<sup>1</sup>, and plastic syringes<sup>2</sup></p> 	<p>examples include: food shrink wrap, packaging that has a plastic cavity or pocket<sup>3</sup></p> 	<p>examples include: food bottles<sup>1</sup>, plastic syringes<sup>2</sup></p> 	<p>examples include: dosing cups or spoons<sup>2</sup>, yogurt containers, bottles, and straws<sup>3</sup></p> 	<p>examples include: vending cups and clamshell carry-out containers<sup>3</sup></p> 	<p>examples include:</p> <ul style="list-style-type: none"> <li>• stainless steel measuring/drinking cups, and containers<sup>6</sup></li> <li>• glass measuring/drinking cups, and containers<sup>2,6</sup></li> <li>• polyurethane feeding tubes and containers<sup>5,6</sup></li> <li>• silicone feeding tubes and containers<sup>2</sup></li> </ul> 





**DISCLAIMER:** Containers are shown as examples only. Containers that fit into example categories may not be made of the listed material, and thus may not be compatible with DOJOLVI. Not all parts of these containers may be compatible with DOJOLVI. Containers may need to be replaced after a period of time due to material wear and tear.

**SOURCES:** 1. Marsh K, Bugusu B. J Food Sci 2007; 72(3):R39-55. 2. Data on File. Ultragenyx Pharmaceutical Inc. Summary Non-clinical Compatibility and Stability Data\_Tubing and Food. 2019. 3. ChemicalSafetyFacts.org Website: [www.chemicalsafetyfacts.org/types-plastic-food-packaging-safety-close-look/](http://www.chemicalsafetyfacts.org/types-plastic-food-packaging-safety-close-look/) 4. Raheem D., Emir J Food Agric. 2013; 25:177-188. 5. S. Sartori, Aliment Pharmacol Ther 2003; 17: 853-856. 6. DOJOLVI (triheptanoin) US Prescribing Information, October 2023. Design adapted from, DOJOLVI Tips for Storing, Mixing and Administering Brochure: PANTHERx Rare Pharmacy of Pittsburgh, PA.

# Materials Compatibility Reference (continued)

**DOJOLVI**<sup>®</sup>  
TRIHEPTANOIN  
Oral Liquid

## Preparation and Administration Instructions

-  Take or give DOJOLVI by mixing well with soft food/liquid (oral administration) or medical food/formula (feeding tube administration). Do not take DOJOLVI alone to avoid stomach upset and breakdown of the feeding tube.
-  Mix or give DOJOLVI using containers, oral syringes, or measuring cups made of materials such as stainless steel, glass, high density polyethylene (HDPE), polypropylene, low density polyethylene, polyurethane, and silicone.
-  After use, rinse materials and containers well with cold water and wipe dry.
-  Do not mix or give DOJOLVI using containers, oral syringes, or measuring cups made of polystyrene or polyvinyl chloride (PVC) plastics.

Regularly monitor the containers, dosing components, or utensils that are in contact with DOJOLVI to ensure they are working properly and not breaking down.

**Please see the DOJOLVI (triheptanoin) full Prescribing Information, including the Patient Information, for additional preparation and administration information.**

ultragenyx